

**25-07-2021 SHIFT-2 CHEMISTRY MEMORY BASE**

1. 3 gm of 'X' dissolve in 100 gm of CCl<sub>4</sub> which increases the boiling point by 0.6. Find molar mass of 'X'. Given K<sub>b</sub> of CCl<sub>4</sub> = 5K kg/mol.

Key:- 250

Sol:-  $\Delta T_b = K_b m$

$$= 5 \times \left( \frac{Wt \times 1000}{M.M \times \text{Mass of solvent (g)}} \right)$$

$$= 5 \times \frac{3 \times 1000}{M.M \times 100}$$

$$0.6 = \frac{150}{M.M}$$

$$M.M = \frac{150}{0.6} = 250 \text{ g/mol.}$$

2. In the following ions, The spin only magnetic moment of of Ti<sup>3+</sup>; Sc<sup>3+</sup>; V<sup>2+</sup> respectively are

- 1) 1.73, 0, 3.87      2) 1.73, 3.87, 0      3) 3.87, 0, 1.73      4) 0, 1.73, 3.87

Key:- 1

Sol:- Ti<sup>3+</sup> {Unpaired electron=1}

Sc<sup>3+</sup> {Unpaired electron=0}

V<sup>2+</sup> {Unpaired electron=3}

$$\mu = \sqrt{n(n+2)} B.M.$$

3. Heat given to a system is 150 joules and work done by the system is 200 joules. The magnitude of the change in the internal energy is:

key:- 50

Sol: q=150 Joules

W=-200 Joules

$$\Delta E = q + w \quad (\text{First law of thermodynamics})$$

$$\Delta E = 150 + (-200)$$

$$\Delta E = -50 \text{ Joule}$$

4. The covalent radii of  $F^- = 1.33 \overset{0}{\text{Å}}$ ,  $O^{2-} = 1.40 \overset{0}{\text{Å}}$  and for  $N = 0.74 \overset{0}{\text{Å}}$ . Then which of the following is incorrect:

- 1) Ionic radius of  $N^{3-}$  is in between  $F^-$  and  $O^{2-}$  but greater than 'N'.
- 2) Ionic radius of  $N^{3-}$  is greater than both  $F^-$  and  $O^{2-}$  and greater than 'N'.
- 3) Ionic radius of  $N^{3-}$  less than both  $F^-$  and  $O^{2-}$  and less than 'N'.
- 4) Ionic radius of  $N^{3-}$  is less than both  $F^-$  and  $O^{2-}$  but less than 'N'.

Key:- 2

Sol:-  $F^- = 1.33 \overset{0}{\text{Å}}$        $N = 0.74 \overset{0}{\text{Å}}$

$O^{2-} = 1.40 \overset{0}{\text{Å}}$        $N^{3-} = 1.46 \overset{0}{\text{Å}}$

Ionic radius of  $N^{3-}$  is greater than  $F^-$  and  $O^{2-}$  size of Anion  $\propto$  Magnitude of -Ve charge

5. For the following ions correct order of their Bond order is:



- 1)  $O_2^- > O_2 > O_2^+ > O_2^{2-}$
- 2)  $O_2^+ > O_2 > O_2^+ > O_2^{2-}$
- 3)  $O_2^+ > O_2 > O_2^- > O_2^{2-}$
- 4)  $O_2^+ > O_2^{2-} > O_2 > O_2^-$

Key:- 2

Sol:- According to Molecular orbital theory

$$B.O = \frac{1}{2} [N_b - N_a]$$

$O_2 = 2.0$

$O_2^+ = 2.5$

$O_2^- = 1.5$

$O_2^{2-} = 1.0$

6.

Colloid	Dispersion Medium
a) Pumice Stone	(i) Liquid in Liquid
b) Cloud	(ii) Gas in solid
c) Cheese	(iii) Liquid in Gas
d) Hair Cream	(iv) Liquid in solid

The correct option is

- 1) a-(iii), b-(ii), c-(i), d-(iv)                      2) a-(i), b-(iv), c-(ii), d-(iii)  
 3) a-(ii), b-(iii), c-(iv), d-(i)                      4) a-(iv), b-(i), c-(iii), d-(ii)

Key:- 3

Sol:-

Dispersed phase	Dispersion medium	Type of colloid	Examples
Solid	Solid	Solid sol	Some coloured glasses and gem stones
Solid	Liquid	Sol	Paints, cell fluids
Solid	Gas	Aerosol	Smoke, dust
Liquid	Solid	Gel	Cheese, butter, jellies
Liquid	Liquid	Emulsion	Milk, hair cream
Liquid	Gas	Aerosol	Fog, mist, cloud, insecticide sprays
Gas	Solid	Solid sol	Pumice stone, foam rubber
Gas	Liquid	Foam	Froth, whipped cream, soap lather

7. In  $Ho^{3+}$  [Atomic No.=67), number of 4f electron are:

Key:- 10

Sol:- Holmium ( $Z=67$ ) :  $4f^{11}, 6s^2$

8.

Column-I	Column-II
a) Li	(i) I is least soluble
b) Na	(ii) Bicarbonate is used in fire extinguisher
c) K	(iii) Carbonate easily decomposed on heating
d) Cs	(iv) Has vital role in biological system

The correct option is:

- 1) a-(iv), b-(iii), c-(ii), d-(i)                      2) a-(iii), b-(ii), c-(iv), d-(i)  
 3) a-(i), b-(iv), c-(ii), d-(iii)                      4) a-(iv), b-(ii), c-(iii), d-(i)

Key:- 2

Sol:- (i)  $Li_2CO_3 \xrightarrow{\Delta} Li_2O + CO_2$


(ii)  $NaHCO_3$  is used in dry fire extinguishers.

(iii) Potassium has vital role in biological systems.

(iv) CsI is least soluble due to smaller hydration enthalpy of its two ions.

9. The concentration of  $H_3O^+$  ions in 0.005 M solution of  $Ba(OH)_2$  at 298 K is  $[x] \times 10^{-12}$ . Assume that  $Ba(OH)_2$  is completely ionized under given conditions.

Key:- 1



Sol:-

0.005M	–	–
–	0.005M	0.010M

$$[OH^-] = 0.01 = 10^{-2} M$$

$$\text{Now } [H^+][OH^-] = K_w$$

$$[H_3O^+] = \frac{K_w}{[OH^-]}$$

$$= \frac{10^{-14}}{10^{-2}} = 10^{-12} M$$

10. Which form interstitial hydride easily?

1) Fe                      2) Cr                      3) Ni                      4) Co

Key:- 2

Sol:- These are formed by many f-block elements. However, the metals of group 7,8 and 9 do not form hydride. Even from group 6, only chromium forms CrH.

11. Match List-I with List-II

Column-I

Column-II

(a) Froth Floatation

(i) Sulphide ore

(b) Bessemer convertor

(ii) Pig iron

(c) Blast furnance

(iii) Ag

(d) Leaching

(iv) Blister copper

1) a-(i), b-(iii), c-(iv), d-(ii)





2) a-(i), b-(iv), c-(ii), d-(iii)

3) a-(iv), b-(ii), c-(iii), d-(i)

4) a-(iii), b-(iv), c-(ii), d-(i)


Key:- 2

12. In which of the following reaction oxidation state changes by 5.




Key:- 2

Sol:-




13. In which of the following compounds one  $\pi$  bond is present and maximum canonical structures possible.



Key:- 2

Sol:-



3 canonical structures

14. An  $e^-$  moving with a velocity of  $2 \times 10^6 m/s$ . If the speed can be measured with an accuracy of 0.02% calculate the uncertainty in its position is  $1.45 \times 10^{-x}$ . The value of x:

Key:- 7

Sol:-  $\Delta x \cdot \Delta p = \frac{h}{4\pi}$

$$\Delta x \cdot m \Delta v = \frac{h}{4\pi}$$

$$\Delta v = 2 \times 10^6 \times \frac{0.02}{100}$$

$$\Delta v \rightarrow 400 m/s$$

$$\Delta x = \frac{h}{4\pi \cdot m \cdot \Delta v} = \frac{6.63 \times 10^{-34}}{4 \times 3.14 \times 9.1 \times 10^{-31} \times 400 m/s} = 1.45 \times 10^{-7} s$$

15.  $A \rightarrow B$

In this reaction, concentration of B changes by 0.2 in 30 minutes. The average rate of the reaction is  $x \times 10^{-1}$  moles per litre hour. The value of x is:

Key:- 4

Sol:- 
$$\text{Rate} = \frac{d[B]}{dt} = \frac{0.2 \text{ mole}^{-1}}{0.5 \text{ hours}}$$

$$\Rightarrow 0.4 \text{ mole} / \text{L hr}$$

16. Which among the following compound is most stable:

- 1)  $[Cr(en)_2(NH_3)_2]Cl_3$                       2)  $[Cr(en)_3]Cl_3$   
 3)  $[Cr(en)(NH_3)_4]Cl_3$                       4)  $[Cr(NH_3)_6]Cl_3$

Key:- 2

Sol:- Chelation due to bidentate ligand. Greater the chelation greater is the stability.

17. In Kjeldahl's method, 0.8 g of organic compound is used. The percentage of Nitrogen came out to be 42%. The \_\_\_\_ ml of 1M  $H_2SO_4$  used to neutralize ammonia.

- 1) 17                      2) 20                      3) 30                      4) 12






Key:- 4

Sol:- 
$$\%N = \frac{1.4N.V}{W}$$

$$42 = \frac{1.4 \times (1 \times 2) \times V}{0.8}$$



$$V = 12 \text{ ml}$$

18. Find the product 'P'.




Key:- 1

Sol:-



Key:- 2

Sol:-




20. Biodegradable polyamide is formed by-

- 1) Glycine + isoprene
- 2) Glycine + Aminocaproic acid
- 3) Alanine + chloroprene
- 4) Acrylonitrile + Aminocaproic acid

Key:- 2

Sol:- Nylon 2-Nylon 6(Polyamide copolymer) is biodegradable polymer.

Its monomer units are : Glycine + Aminocaproic acid




21. Benzenenitrile with Grignard reagent form product (P), which of the following chemical test given by product (P)



- 1) Schiff's reagent
- 2) Iodoform
- 3) Ninhydrin
- 4) Tollen's test

Key:- 2

Sol:-



22. Correct order of acidic strength form following compounds:



1) a>b>c>d

2) d>c>d>a





3) b>c>d>a

4) c>b>a>d

Key:- 2

Sol:- Acidic strength  $\propto$  stability of conjugate base.




23. Structure of cytosine is:





Ans 1

Sol. NCERT

24.








Ans 3

Sol.



25. S<sub>1</sub>: CFCs are dissociated with to Cl radical by radiation of visible region.

S<sub>2</sub>: O<sub>3</sub> reacts with nitric oxide to form N<sub>2</sub> & O<sub>2</sub>

1) False, True

2) False, False

3) True, False

4) True, True

Ans 2

Sol:  $CFCs + UV \rightarrow Cl^*$

$O_3 + NO \rightarrow NO_2 + O_2$

26. Increasing order of density:

(I) Benzene

(II) 1,3-Dichlorobenzene

(III) Chloro benzene

(IV) 1-Bromo-3-chlorobenzene

1) IV > II > III > I

2) IV > III > II > I

3) III > II > IV > I

4) I > II > III > IV

Ans 1

Sol: Higher the molecular weight higher will be density.

27. Maleic anhydride can be prepared by

1) Treating cis but-2-ene-1,4-dioic acid with alcohol.

2) Heating cis but-2-ene,1,4-dioic acid

3) Treating trans but-2-ene-1,4 dioic acid with alcohol and acid

4) Heating trans but-2-ene-idric acid

Ans 2

Sol.

